

Reducing violence against women through psychological interventions targeting men

Men and boys are the main perpetrators of violence against women.¹

Psychological issues stemming from mental health problems or prolonged conflict are risk factors associated with men perpetrating violence.^{2,3,4} Moreover, patriarchy and gender norms often normalise violence against women and girls. To successfully reduce violence against women, prevention interventions need to include men and boys.⁵

The Sexual Violence Research Initiative (SVRI), in partnership with the World Bank Group, has funded several studies in East and Central Africa that target men. These interventions attempt to address local gender norms and context-specific risk factors associated with men abusing women.

ADDRESSING VIOLENCE IN A CONFLICT-AFFECTED SETTING

Years of conflict in the Democratic Republic of Congo have had serious psychological impacts on both men and women. A 2012 study⁶ undertaken



by Promundo revealed links between men's exposure to trauma and violence and their use of violence against partners. The study showed how men often cope with psychological problems and distress by using strategies that affirm social expectations that men should dominate women. These strategies include alcohol use, sexual promiscuity, physical or psychological

violence, and rejecting partners who have been raped.

In response to these findings, in 2013 Promundo piloted "Living Peace",⁷ a 15-week intervention targeting 324 men and their partners. Living Peace helps men and their partners to develop positive coping strategies and restore violence-free relationships in post-conflict and high-violence settings. During the intervention, men discussed how war and conflict influenced their idea of what it means to "be a man", explored their trauma, and learnt how to cope differently with stress, loss and anger. Through these discussions and educational activities men learnt to adopt non-violent coping strategies.

Three years later, Promundo conducted a follow-up study⁸ that found a lasting reduction in intimate partner violence. The changes in men's behaviour and attitudes have positively affected the lives of women and children, extending to the broader community.

www.svri.org svri@mrc.ac.za

#GBVSolutions @WBG_Gender @TheSVRI



The Sexual Violence Research Initiative is hosted by the South African Medical Research Council

MENTAL HEALTH AND INTIMATE PARTNER VIOLENCE

Recognising gendered coping strategies and the links between men with common mental health problems (such as depression, anxiety and post-traumatic stress disorder), alcohol and substance use, and high incidences of intimate partner violence, World Vision Kenya is assessing whether a mental health intervention for men may help them better manage stressors associated with their use of violence against their partners.⁹

This study is adapting the World Health Organization's Problem Management

THE PROGRAMME

The Problem Management Plus programme¹⁰ is a psychological intervention targeting individuals with emotional distress. It comprises five sessions that deal with problem management and behavioural strategies.

Plus programme, which was offered to women affected by violence in peri-urban communities in Nairobi, to a group format for men with common mental health problems. The intervention will also raise awareness about intimate partner violence within the community through, for example, community dialogues.

INVOLVING MEN IN VIOLENCE PREVENTION INTERVENTIONS

Although numerous interventions have been designed to involve men and boys in violence prevention activities, many of these have struggled to effectively engage them. In response to this problem, the Rakai Health Sciences Program and the University of California, San Diego, are working to create an evidence-based framework for effectively engaging men and boys in programmes that aim to transform gender norms and reduce violence against women and girls.

This study will build on the "Safe Homes and Respect for Everyone" (SHARE) project, developed by the

Rakai Health Sciences Program. SHARE was the first behavioural intervention to significantly reduce HIV incidence and community-wide levels of intimate partner violence.¹¹ Drawing on lessons learnt during the implementation of SHARE, the project team will assess gender norms, and identify factors leading to men perpetrating violence against their intimate partners and barriers to their engagement in prevention programmes.¹²

Results from this study will be used to inform the delivery and expansion of an intervention to prevent the perpetration of intimate partner violence in sub-Saharan Africa and other low- and middle-income settings.

WAY FORWARD

The SVRI and World Bank Group support teams in widely disseminating these research findings to ensure that evidence from these studies informs policy and practice. ■

¹ Krug, E.G., Dahlberg, L.L., Mercy, J.A., Zwi, A.B. & Lozano, R. Eds. 2002. *World report on violence and health*. Geneva: World Health Organization.

² Campbell, J.C. 2002. Health consequences of intimate partner violence. *Lancet*. 359(9314): 1331-1336.

³ Campbell, J.C. & Soeken, K. 1999. Forced sex and intimate partner violence: Effects on women's health. *Violence Against Women*. 5(9): 1017-1035.

⁴ Ellsberg, M., Jansen, H.A., Heise, L., Watts, C.H., Garcia-Moreno, C. & WHO multi-country study on women's health and domestic violence against women study team. 2008. Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: An observational study. *Lancet*. 371(9619): 1165-1172. DOI: 10.1016/S0140-6736(08)60522-X.

⁵ Jewkes, R., Flood, M. & Lang, J. 2015. From work with men and boys to changes of social norms and reduction of inequities in gender relations: A conceptual shift in prevention of violence against women and girls. *Lancet*. 385(9977): 1580-1599. DOI: 10.1016/S0140-6736(14)61683-4.

⁶ Slegel, H., Barker, G. & Levtov, R. 2014. *Gender relations, sexual and gender-based violence and the effects of conflict on women and men in north Kivu, eastern Democratic Republic of the Congo: Results from the International Men and Gender Equality Survey*. Washington, D.C., and Cape Town, South Africa: Promundo-US & Sonke Gender Justice. Available: <http://promundoglobal.org/2014/06/06/images-drc/> [2017, May 30].

⁷ Promundo-US. 2014. Living Peace groups implementation manual and final project report: GBV prevention and social restoration in the DRC and Burundi. *LOGiCA Study Series*. Available: <http://promundoglobal.org/wp-content/uploads/2014/12/Living-Peace-Groups-Implementation-Manual-and-Final-Project-Report.pdf> [2017, June 2].

⁸ Slegel, H. & Tankink, M. 2016. *When no one is looking: Three years of healing in Democratic Republic of the Congo*. Available: <http://www.svri.org/blog/when-no-one-looking-3-years-healing-democratic-republic-congo> [2017, April 12].

⁹ Koyiet, P.N. 2016. *Mental health and intimate partner violence in Kenya*. Available: <http://www.svri.org/blog/mental-health-and-intimate-partner-violence-kenya> [2017, April 12].

¹⁰ World Health Organization. 2016. *Problem Management Plus (PM+): Individual psychological help for adults impaired by distress in communities exposed to adversity*. Generic field-trial version 1.0. Geneva: World Health Organization.

¹¹ Wagman, J.A., Gray, R.H., Campbell, J.C., Thoma, M., Ndyababo, A., Ssekasanvu, J., Nalugoda, F., Kagaayi, J., Nakigozi, G., Serwadda, D. & Brahmbhatt, H. 2015. Effectiveness of an integrated intimate partner violence and HIV prevention intervention in Rakai, Uganda: Analysis of an intervention in an existing cluster randomised cohort. *Lancet Global Health*. 3(1): e23-33. DOI: [http://dx.doi.org/10.1016/S2214-109X\(14\)70344-4](http://dx.doi.org/10.1016/S2214-109X(14)70344-4).

¹² Bonnevie, E., Wagman, J., Nalugoda, F., Nakigozi, G., Nakyanjo, N. & Fleming, P. 2016. *Engaging men and boys in the prevention of violence against women in Uganda*. Available: <http://www.svri.org/blog/engaging-men-and-boys-prevention-violence-against-women-uganda> [2017, April 12].

This research was funded through the SVRI Grant and the SVRI and World Bank Group Development Marketplace. For more information on the grants, see <http://www.svri.org/what-we-do/research-support/svri-grant>.